

County Surveyor's Record, Douglas County, Oregon

Dependent Resurvey of a Portion of the Boundaries
of Donation Land Claim No. 47,
T. 28 S., R. 7 W., Willamette Meridian, Oregon

15423

CHAINS

- 19.70 Creek, 1 lk. wide, course S. 85° E.; asc. 30 ft. over SE. slope.
- 21.65 Spur, slopes E.; desc. 2 ft. over NE. slope.
- 21.85 Ravine, drains S. 75° E., asc. 140 ft. over SE. slope.
- 32.15 Ravine, drains S. 65° E.; asc. 105 ft. over SE. slope.
- 37.80 Ridge, bears N. 80° W. and S. 80° E.; desc. 115 ft. over NE. slope.
- 43.65 Ravine, drains E.; asc. 20 ft. over SE. slope.
- 57.31 The NW. cor. of Claim No. 47, on the N. bdy. of sec. 15, hereinbefore described.
From this point, the cor. of secs. 9, 10, 15, and 16, bears N. 88. 35' W., 13.22 chs. dist.
Land, mountainous.
Soil, rocky clay.
Timber, fir, hemlock, cedar, maple, oak, myrtle, alder, and chinquapin; undergrowth, salal, vine maple, Oregon grape rhododendron, poison oak, and fern.

GENERAL DESCRIPTION

The lands included in the notes of the foregoing resurvey are located in mountainous terrain about 2 1/2 miles southwesterly from the hamlet of Lookingglass, Oregon. The area is drained by Olalla Creek, and many of its tributaries. The elevations range from about 640 feet above sea level near the corner of sections 10, 11, 14, 15, to about 1,650 feet above sea level near the corner of sections 15, 16, 21, and 22.

Access into the area is by way of a gravelled road, which branches from Lookingglass Road.

Timber in the area consists of fir, hemlock, cedar, maple, oak, alder, myrtle, and chinquapin; with undergrowth of salal, poison oak, vine maple, rhododendron, Oregon grap, and fern.

The soil is composed of rocky clay loam, from a few inches to several feet in depth, overlaying a sandstone bedrock which is exposed in numerous outcroppings throughout the area.

No evidence of mineral activity was noted in the area covered by this resurvey.

The average of a considerable number of readings over all parts of the area gives a value of 18° 30' E., for the mean magnetic declination. There is a range of up to 0° 15' in local attraction.