

## County Surveyor's Record, Douglas County, Oregon

Dependent Resurvey, Subdivisional Lines, T. 25 S., R. 7 W.

## CHAINS

Set an iron post, 28 ins. long,  $2\frac{1}{2}$  ins. diam., 24 ins. in the ground, with brass cap mkd.

W  $\frac{1}{16}$  S 18  
1964

from which

A fir, 12 ins. diam., bears S.  $79\frac{1}{2}^{\circ}$  E., 43 lks. dist., mkd. W  $\frac{1}{16}$  S18 BT.

A white fir, 17 ins. diam., bears S.  $38\frac{1}{4}^{\circ}$  W.,  $44\frac{1}{2}$  lks. dist., mkd. W  $\frac{1}{16}$  S18 BT.

- 60.20 Descend 15 ft. over NW. slope.  
Ravine, drains N.; asc. 160 ft. over E. slope.
- 66.00 Top of ascent; slopes N.; desc. 10 ft. along N. slope.
- 71.00 Ravine, drains N.  $30^{\circ}$  E.; asc. 80 ft. over E. slope.
- 73.00 Spur, slopes N.  $20^{\circ}$  W.; desc. 40 ft. over W. slope.
- 77.09 Intersect the W. bdy. of the Tp. at the closing corner of secs. 7 and 18, previously described.
- From this point the cor. of secs. 12 and 13, on the E. bdy. of T. 25 S., R. 8 W., bears N.  $0^{\circ} 15'$  E., 9.93 chs. dist., previously described.

Land, mountainous.  
Soil, rocky clay loam  
Timber, fir, hemlock, cedar, white fir, madrone, alder, dogwood, yew, oak, maple, and myrtle; undergrowth, vine maple, salal, ocean spray, hazel, Oregon grape, poison oak, manzanita, rhododendron, huckleberry, & blackberry.

The lands included in the foregoing survey are situated about 10 miles westerly of the hamlet of Umpqua, Oregon. Access is gained by way of Hubbard Creek Road and Tyee Road, and by several minor roads which branch off from them.

The area is drained by the Umpqua River and some of its tributaries of which Hubbard Creek, Cougar Creek, and Little Wolf Creek are the most prominent. The area is rough and mountainous, elevations range from about 300 feet above sea level at the Umpqua River, in section 6, to about 1,995 feet above sea level on the ridge in section 19. Soil is composed generally of rocky clay loam with some sandy clay loam along the river bottom land. There were several small farms and orchards noted along Hubbard Creek and along the Umpqua River.

Timber consists primarily of fir, with stands of hemlock, cedar, white fir, madrone, alder, dogwood, yew, maple, and myrtle. The undergrowth includes vine maple, salal, ocean spray, hazel, Oregon grape, poison oak, manzanita, rhododendron, huckleberry, and blackberry.

No mineral deposits were noted in the area.

The average of a considerable number of readings throughout the area gives a value of  $19^{\circ} 24'$  E. for the mean magnetic declination, with a range of  $0^{\circ} 45'$  in local attraction.