

County Surveyor's Record, Douglas County, Oregon

Dependent Resurvey, Portion of Subdivisional Lines, T. 26 S., R. 7 W.

CHAINS

A fir, 9 ins. diam., bears N. 55° E., 186 lks. dist.,
mkd. E 1/16 S8 BT.

A fir, 20 ins. diam., bears S. 6° E., 96 lks. dist.,
mkd. E 1/16 S17 BT.

The corner falls at the SE. edge of rocky barren.

Descend 215 ft. over broken SW. slope.

39.36

The $\frac{1}{4}$ sec. cor. of secs. 8 and 17, perpetuated and recorded by the County Surveyors Office in 1966, monumented with an Iron post, $1\frac{1}{2}$ ins. diam., protruding 12 ins. above the ground, firmly set, with brass cap mkd.

COUNTY SURVEYORS OFFICE

T 26 S
 $\frac{1}{4}$ $\frac{S 8}{S 17}$
R 7 W
1966

from which the remaining original bearing tree

A cedar, 44 ins. diam., bears N. 80° W., 21 lks. dist.,
with original scribe marks $\frac{1}{4}$ S BT visible on bark.
This tree was rescribed $\frac{1}{4}$ S8 CS BT on bark in 1966.

and bearing tree mkd. by the County Surveyors Office in 1966.

A fir, 13 ins. diam., bears N. 57° E., 6 lks. dist.,
with illegible scribe marks visible on partially
healed double blaze.

A fir, 16 ins. diam., bears S. 7° W., 27 lks. dist.,
with illegible scribe marks visible on partially
healed double blaze.

Land, mountainous.

Soil, rocky clay with rock outcroppings.

Timber, fir, hemlock, cedar, alder, maple, chinquapin, and
yew; undergrowth, young timber, fern, salal, hazel, arrowwood,
poison oak, huckleberry, and Oregon grape.

GENERAL DESCRIPTION

The portion of township 26 south, range 7 west, covered by this resurvey is located in the Coast Range of mountains about 12 miles outwest of Sutherlin, Oregon. Elevations along the lines resurveyed range from about 630 feet above sea level, near the cor. of secs. 3, 4, 9 and 10, to about 2400 feet above sea level, on Bear Ridge on the line between secs. 9 and 16. The soil is a rocky clay with rock outcroppings and suited mainly for the growing of timber. The timber is predominantly fir with a scattering of hemlock, cedar, alder, madrone, maple, yew, chinquapin, and oak. Undergrowth consists chiefly of young timber, fern, salal, poison oak, Oregon grape, hazel, arrowwood, vine maple, huckleberry, blackberry, salmonberry, and ceanothus.

The area is drained by tributaries of Hubbard Creek and Mill Creek, both of which drain into the Umpqua River. Several access roads lead into the area from hard surfaced roads along these creeks to old logging operations. No permanent improvements were noted in the area.

No mineral deposits were noted in the area.

The average of a number of readings along the lines resurveyed gives a value of 19° 10' E. for a mean magnetic declination. There is a range of 1° in local attraction.