

County Surveyor's Record, Douglas County, Oregon

14889

Dependent Resurvey, Portion of Subdivisional Lines
T. 24 S., R. 4 W., Willamette Meridian, Oregon

CHAINS

- 37.00 Spur, slopes SE.; desc. 40 ft. over NW. slope.
- 39.68 The 1/4 sec. cor. of secs. 27 and 34, perpetuated and recorded by Milo E. Godfrey, Deputy County Surveyor, in 1963, monumented with an iron pipe, 2 ins. diam., firmly set, projecting 6 ins. above the ground, with brass cap mkd.

COUNTY SURVEYORS OFFICE

T 24 S
1/4 $\frac{S\ 27}{S\ 34}$
R 4 W
1963

from which the remains of the original bearing trees

A stump hole, bears N. 20° E., 90 lks. dist.

A madrone root crown, size indeterminate
bears S. 76° W., 31 lks. dist.

and bearing trees marked by Godfrey

A fir, 15 ins. diam., bears N. 31° E.,
33 lks. dist., with healed blaze.

A fir, 9 ins. diam., bears S. 35° W.,
23 lks. dist., with healed blaze.

Land, ountainous.

Soil, rocky clay loam.

Timber, fir, hemlock, alder, madrone, white oak, black oak, cedar, dogwood, and maple; undergrowth, salal, hazel, fern, poison oak, and vine maple.

GENERAL DESCRIPTION

The lands included in the foregoing survey are situated in Douglas County, about 8 miles northeasterly of the town of Sutherlin, Oregon. Access is by way of a paved county road which leads from Sutherlin, Oregon to Nonpareil, Oregon, and several logging roads which branch off from it.

The area is drained by several small creeks draining into Long Valley Creek and Contell Creek, which flow in a general southwesterly direction. Elevations range from about 960 feet above sea level, near the corner of sections 26, 27, 34 and 35, to about 1400 feet above sea level, near the 1/4 section corner of sections 26 and 35.

The soil is composed generally of rocky clay loam. and is unsuited for agricultural purposes, but supports a luxuriant growth of timber, which is primarily fir and alder, with stands of hemlock, madrone, maple, white oak, and black oak. The undergrowth consists chiefly of salal, vine maple, hazel; poison oak, and fern.'

No evidence of mineral deposits were noted along the lines surveyed.

The average of a considerable number of readings throughout the area resurveyed gives a value of 19° 30' E., for the mean magnetic declination. There is a range of up to 0° 30' in local attraction.