

County Surveyor's Record, Douglas County, Oregon

15298

Dependent Resurvey, Portion of Subdivisional Lines,
T. 24 S., R. 7 W., Willamette Meridian, Oregon

CHAINS

79.72

A fir, 12 ins. diam., bears N. 68° W., 23 lks.
dist., mkd. X BT.

A fir, 18 ins. diam., bears S. 39° W., 33 lks.
dist., mkd. X BT.

Ascend 200 ft. over SE. slope.

The cor. of secs. 28, 29, 32, and 33, perpetuated by James W. Byron, Deputy County Surveyor, in 1960, subsequently perpetuated and recorded by Bruce Shaner, Registered Engineer No. 4906, in 1972, monumented with an iron post, 1 1/2 ins. diam., firmly set, protruding 10 ins. above ground, with brass cap mkd.

COUNTY SURVEYORS OFFICE

T 24 S R 7 W

S 29 | S 28

S 32 | S 33

1960

from which the remains of the original bearing trees

An oak, 32 ins. diam., bears N. 3° E., 255 lks.
dist., no marks visible on decayed opened blaze.

and bearing trees mkd. by Byron, in 1960

A white oak, 22 ins. diam., bears N. 59° E., 82 lks.
dist., with healed blaze.

A white oak, 20 ins. diam., bears S. 67 1/2° E., 97
lks. dist., with healed blaze.

A white oak, 18 ins. diam., bears N. 81° W., 35 1/2
lks. dist., with healed blaze.

and a bearing tree mkd. by Shaner, in 1972

A fir, 10 ins. diam., bears S. 62° W., 25 lks.
dist., with scribe marks T24S R7W S32 BT visible
on opened blaze.

Land, mountainous.

Soil, rocky clay loam.

Timber, fir, cedar, hemlock, white oak, maple, madrone,
chinquapin, and alder; undergrowth, vine maple, salal,
fern, salmonberry, blackberry, hazel, rhododendron, and
poison oak.

GENERAL DESCRIPTION

The lands included in the notes of the foregoing survey are located in Douglas County, about 15 miles northwest of the town of Sutherlin, Oregon. The area is reached by way of Oregon State Highway Number 225, from Sutherlin.

The elevations in the area range from about 200 feet above sea level through the bottom land along the Umpqua River to about 1,850 feet above sea level near the southeast corner of section 33. The soil is rocky clay loam. The timber stands are primarily composed of fir, hemlock, and cedar, with lesser amounts of alder, madrone, yew, chinquapin, maple, and oak scattered throughout the area. The undergrowth is generally dense and consists of vine maple, hazel, fern, salal, salmonberry, blackberry, ocean spray, ceanothus, and poison oak.