

Dependent Resurvey, Portion of Subdivisional Lines,
T. 27 S., R. 9 W., Willamette Meridian, Oregon

CHAINS

From this corner, the original closing corner, bears N. 88° 47' E., 0.22 chs. dist., monumented with the broken remains of the original sandstone, size indeterminate, loosely set 4 ins. in the ground, with 4 grooves and CC visible on broken pieces. I add the marks AM and bury the remains of the stone below the surface of the ground, from which the original bearing trees:

A fir, 44 ins. diam., bears S. 51° W., 27 lks. dist., with illegible scribe marks visible on opened blaze.

A fir, 47 ins. diam., bears N. 85° W., 93 lks. dist. with healed blaze.

From this same corner, the cor. of secs. 19 and 30 only, on the W. bdy. of T. 27 S., R. 8 W., bears N. 1° 00' W., 1.54 chs. dist., hereinbefore described.

Land, mountainous.

Soil, rocky clay loam.

Timber, fir, hemlock, cedar, yew, and chinquapin; undergrowth, vine maple, rhododendron, sword fern, and hazel.

GENERAL DESCRIPTION

The lands included in the foregoing survey are located in mountainous terrain about 12 miles northerly of Camas Valley, Oregon. The area is located on a divide and is drained by tributaries of Williams River to the north and tributaries of the East Fork Coquille River to the south. The elevation ranges from about 2,100 feet above sea level near the $\frac{1}{4}$ section corner of section 30 only, to about 3,000 feet above sea level near the corner of sections 19 and 30 on the west boundary of township 27 south, range 8 west.

Access into the area is by way of Burnt Mountain access road and Burnt Mountain Ridge Road, both oiled roads which traverse the main ridges into the area and converge in the immediate area, via Middle Creek Road and the Coos Bay Wagon Road.

Soil in the area is composed of sandy clay loam from a few inches to several feet in depth, overlaying sandstone bedrock. The lands are forested with fir, hemlock, cedar, yew, and chinquapin; the undergrowth consists of vine maple, rhododendron, sword fern, and hazel.

No evidence of mineral activity was noted in the area covered by the resurvey.

The average of a considerable number of readings throughout the area gives a value of 18° 45' E. for the mean magnetic declination. There is a range of 0° 15' due to local attraction.