

Subdivision of Section 17, T. 24 S., R. 7 W.,
Willamette Meridian, Oregon

CHAINS	
	N. 89° 44' W., on the E. and W. center line of the NW $\frac{1}{4}$ of sec. 17.
	Ascend 290 ft. over NE. slope, through open field.
3.20	From this point, the NE. corner of a wood frame barn, 52 x 40 ft., bears South, 0.27 chs. dist., long side bears E. and W.
12.30	Dirt road, 30 lks. wide, bears N. 55° W. and S. 55° E.
17.40	Ridge, bears N. 65° W. and S. 65° E.; desc. 55 ft. over SW. slope.
19.965	The NW 1/16 sec. cor. of sec. 17. Descend 130 ft. over SW. slope.
25.80	Gravelled road, 40 lks. wide, bears N. 60° W. and S. 60° E.
26.90	Creek, 7 lks. wide, course S. 40° E.; asc. 155 ft. over NE. slope.
32.65	Spur, slopes N. 15° E.; desc. 15 ft. over NW. slope.
34.90	Creek, 3 lks. wide, course N. 35° E.; asc. 140 ft. over SE. slope.
40.84	The N 1/16 sec. cor. of secs. 17 and 18.
	<p style="text-align: center;">GENERAL DESCRIPTION</p> <p>The lands embraced in this survey are located in mountainous terrain about 12 miles northwesterly from Southerlin, Oregon. The area drains southerly via Basin Creek into the Umpqua River. The elevations range from about 280 feet above sea level at the corner of sections 16, 17, 20, and 21, to about 1,240 feet above sea level at the corner of sections 8, 9, 16, and 17.</p> <p>Access into the area is by way of the Basin Creek access road.</p> <p>Timber in the area consists of fir, hemlock, cedar, oak, maple, madrone, myrtle, alder, and chinquapin; with undergrowth of salal, poison oak, vine maple, rhododendron, Oregon grape, and fern. The soil is composed of rocky clay loam.</p> <p>No evidence of mineral activity was noted in the area covered by this survey.</p> <p>The average of a considerable number of readings over all parts of the area gives a value of 18° 30' E. for the mean magnetic declination. There is a range of up to 0° 45' in local attraction.</p>