

16239

Corrective Dependent Resurvey of a
Portion of the Subdivisional Lines,
T. 32 S., R. 7 W., Willamette Meridian, Oregon

CHAINS

T 32 S R 7 W
S 14 | S 13
S 23 | S 24
1976

from which bearing trees mkd. by Popken and Carlson

A fir, 21 ins. diam., bears N. 33° E., 73 lks.
dist., with scribe marks T32S R7W S13 BT
visible on partially healed blaze.

A fir, 18 ins. diam., bears S. 60½° E., 28 lks.
dist., with scribe marks T32S R7W S24 BT
visible on partially healed blaze.

A fir, 9 ins. diam., bears S. 10° W., 58 lks.
dist., with scribe marks T32S R7W S23 BT
visible on partially healed blaze.

A fir, 19 ins. diam., bears N. 29° W., 135 lks.
dist., with scribe marks T32S R7W S14 BT
visible on partially healed blaze.

Corner is located at the NW. cor. of a recently logged
area in sec. 24.

Land, mountainous.

Soil, rocky clay, with rock outcroppings.

Timber, fir, pine, cedar, oak, madrone, and chinquapin;
undergrowth, manzanita, hazel, arrowwood, poison oak,
fern, and young timber.

GENERAL DESCRIPTION

The lands included in this corrective resurvey are located approximately five miles northwest of the small town of Glendale, Oregon. The area is very mountainous and broken. The elevations range from about 2,200 feet above sea level near the corner of sections 19, 24, 25, and 30, to about 4,000 feet above sea level near the summit of Crayback Mountain which is situated in the SE¼ of section 13. The divide between Cow Creek and Middle Creek drainages extends diagonally from northeast to southwest, separating the area surveyed nearly equally.

Access into the area is by way of McCollough and Totten Creek Roads.

The soil is rocky clay loam, except on the south slope of Grayback Mountain which is mostly loose rock and rock outcroppings. Timber covering this locality ranges from heavy to moderate, with stands of fir, pine, live oak, madrone, chinquapin, and cedar. The undergrowth is generally dense and consists of arrowwood, hazel, poison oak, fern, and young timber.

There is no development in the area nor evidence of having been any in the past.

There was no evidence of mineral activity noted along the lines resurveyed.

The average of a considerable number of readings throughout the area gives a value of 19° 30' E. for the mean magnetic declination, with a range of 1° 30' due to local attraction.