

County Surveyor's Record, Douglas County, Oregon

17237

Dependent Resurvey, Portion of Subdivisional Lines,
T. 21 S., R. 7 W., Willamette Meridian, Oregon

CHAINS

COUNTY SURVEYORS OFFICE

T 21 S
S 27 | S 26
S 34 | S 35
R 7 W
1957

from which the remains of three original bearing trees

A fir stump, 40 ins. diam., bears S. 41° E., 57 lks. dist., with no marks visible.

A yew, 19 ins. diam., bears S. 71° W., 32 lks. dist., with no scribe marks visible on opened blaze. (Record, S. 66° W.)

A fir, 48 ins. diam., bears N. 61 1/2° W., 35 lks. dist., with no marks visible. (Record, N. 65° W.)

and bearing trees mkd. by Byron

A hemlock, 20 ins. diam., bears N. 34° E., 67 1/2 lks. dist., with healed double blaze.

A fir, 28 ins. diam., bears S. 85° E., 67 lks. dist., with healed double blaze.

A hemlock, 24 ins. diam., bears S. 17° W., 61 lks. dist., with healed double blaze.

A fir, 15 ins. diam., bears N. 13 1/2° W., 16 lks. dist., with healed double blaze.

Land, mountainous.
Soil, rocky clay.
Timber, ash, oak, fir, cedar, yew, hemlock, alder, madrone, chinquapin, and maple; undergrowth, poison oak, blackberry, huckleberry, salmonberry, arrowwood, hazel, fern, Oregon grape, salal, vine maple, and young timber.

GENERAL DESCRIPTION

The area covered by this resurvey is located about seven miles northeast of the city of Elkton, Oregon. Access to the area is gained via Tom Folley Creek Road and other gravelled and dirt roads. The elevation varies from about 1440 feet above sea level near the quarter section corner of sections 26 and 35, to about 220 feet above sea level near the corner of sections 1, 2, 35, and 36.

The area is drained by Tom Folley Creek, North Fork Tom Folley Creek, and other small drainages flowing into Tom Folley Creek, which flows southwesterly to Elk Creek.

The soil is mostly rocky clay with some sandy loam in the creek bottoms. The land is used primarily for timber. The timber consists of fir, oak, madrone, hemlock, cedar, yew, chinquapin, and alder. The undergrowth consists of poison oak, vine maple, hazel, arrowwood, salmonberry, huckleberry, salal, Oregon grape, blackberry, cascara, fern, hawthorne, scotch broom, willow, and young timber.

No mineral deposits were noted during the survey.

The average of a number of readings along the lines surveyed gives a mean magnetic declination of 21° E., with a range of 0° 30' in local attraction.