

County Surveyor's Record, Douglas County, Oregon

Dependent Resurvey, Portion of Subdivisional Lines, 17596
T. 22 S., R. 5 W., Willamette Meridian, Oregon

CHAINS

and bearing trees mkd. by McGillicuddy

A fir, 26 ins. diam., bears N. 65° E., 49 lks. dist., with healed blaze.

A fir, 18 ins. diam., bears S. 55° E., 25 lks. dist., with healed blaze.

At the corner point

Set an iron post, 28 ins. long, 2 1/2 ins. diam., 24 ins. in the ground, with brass cap mkd.

T 21 S	R 5 W
S 35	S 36
S 2	S 1
T 22 S	R 5 W

1982

from which the new bearing trees

A grand fir, 6 ins. diam., bears S. 47° W., 44 lks. dist., mkd. T22S R5W S2 BT.

A fir, 16 ins. diam., bears N. 27° W., 21 1/2 lks. dist., mkd. T21S R5W S35 BT.

Land, mountainous.

Soil, rocky clay.

Timber, fir, cedar, chinquapin, yew, hemlock, maple, oak, madrone, and alder; undergrowth, poison oak, vine maple, arrowwood, hazel, Oregon grape, salal, blackberry, fern, wild rose, and young timber.

GENERAL DESCRIPTION

The lands included in the foregoing resurvey are located in mountainous country about five miles east of Drain, Oregon. Access is gained via dirt roads extending into the area.

The area drains northerly into Pass Creek and southerly into Wehmeyer Creek and Salt Creek. The elevations range from about 1,280 feet above sea level near the 1/4 sec. cor. of sections 1 and 12 to about 560 feet above sea level at the corner of sections 1, 6, 7, and 12.

The soil is made up of a rocky clay loam. The timber consists of fir, yew, hemlock, cedar, chinquapin, oak, madrone, cascara, ash, and alder. Undergrowth is mainly hazel, poison oak, Oregon grape, salal, vine maple, blackberry, huckleberry, elderberry, salmonberry, fern, and young timber.

There were no mineral deposits noted along the lines resurveyed.

The average of a number of readings throughout the area gives a value of 19° E. for the magnetic declination. There is a range of 1° due to local attraction.