

# County Surveyor's Record, Douglas County, Oregon

18423

Dependent Resurvey of a Portion of the Subdivisional Lines,  
T. 25 S., R. 1 W., Willamette Meridian, Oregon

## CHAINS

from which the remains of the bearing trees mkd. by Bruce A. Shaner, Registered Engineer No. 4906, in 1967

A sawed fir stump, 26 ins. diam., bears S.  $19\frac{1}{2}^{\circ}$  W.,  
52 $\frac{1}{2}$  lks. dist., with scribe marks  $\frac{1}{4}$  S9 BT visible  
on open blaze.

A sawed fir stump, 26 ins. diam., bears N.  $49^{\circ}$  W.,  
22 $\frac{1}{2}$  lks. dist., with healed blaze.

From this point, a point established and recorded as the  $\frac{1}{4}$  sec.  
cor. of secs. 4 and 9, by Thomas E. Shrum, Jr., Registered  
Engineer No. 4287, in 1966, reported by Jelley, bears  
N.  $81^{\circ} 15' 00''$  W., 1.000 chs. dist., monumented with an iron  
post, 1 in diam., firmly set flush with the ground, with a  
2 $\frac{1}{2}$  in. diam., aluminum cap mkd. USPC PROP COR T25S RIW  $\frac{1}{4}$  S4 S9 RE  
4287, from which the remains of bearing trees mkd. by Shrum:

A sawed cedar stump, 36 ins. diam., bears N.  $31^{\circ}$  E.,  
14 lks. dist., with scribe marks S4  $\frac{1}{4}$  BT visible on  
open blaze.

A chinquapin, 10 ins. diam., bears S.  $45^{\circ}$  W.,  
30 $\frac{1}{2}$  lks. dist., with scribe marks S9  $\frac{1}{4}$  BT  
visible on open blaze.

A chinquapin, 11 ins. diam., bears S.  $41^{\circ}$  E.,  
21 lks. dist., with scribe marks  $\frac{1}{4}$  S9 CS BT  
visible on partially healed blaze.

Land, mountainous.

Soil, rocky clay.

Timber, fir, madrone, cedar, hemlock, yew, alder, chinquapin,  
maple, and dogwood; undergrowth, salal, Oregon grape, manzanita  
ceanothus, arrowwood, hazel, vine maple, blackberry, salmonberry,  
thimbleberry, huckleberry, ferns, and young timber.

## GENERAL DESCRIPTION

The lands included in the foregoing resurvey are locate in  
mountainous country about 18 miles northeasterly of Glide,  
Oregon. The elevation ranges from about 1800 feet above sea  
level at the corner of sections 2, 3, 10, and 11, to about 3000  
feet above sea level near the corenr of sections 3, 4, 9, and 10.

The area is drained by Trapper Creek, Wolverine Creek, and  
other smaller and seasonal drainages flowing easterly into Canton  
Creek. The lands are forested with fir, madrone, alder, pine,  
yew, maple, chinquapin, hemlock, and dogwood. Undergrowth  
consists of vine maple, hazel, arrowwood, blackberry, salmonberry,  
thimbleberry, huckleberry, Oregon grape, salal, manzanita,  
ceanothus, ferns, and young timber. Soil consists of a rocky  
clay.

Access is gained by way of gravelled roads branching from  
Canton Creek Road.

The average of a number of readings throughout the area  
gives a value of  $17^{\circ} 30'$  E., for the mean magnetic declination.  
There is a range of  $1^{\circ}$  of variation due to local attraction.